Surveillance Week 18 (April 27 - May 3, 2025)



Chicago Respiratory Virus Weekly Surveillance Report



Brandon Johnson, Mayor

May 9, 2025

Olusimbo Ige, MD, MS, MPH, Commissioner


This report summarizes key respiratory virus surveillance indicators. The indicators are compiled from laboratory-based data as well emergency department visit data. All data are preliminary and may change as additional reports are received. Historical and seasonal summary reports can be found here: Chicago <u>Influenza and Respiratory Virus Surveillance Report</u>.

A new Respiratory Illness Dashboard is now available on the <u>CDPH website</u>. This dashboard summarizes information about respiratory virus disease activity in Chicago, with a focus on COVID-19 (caused by the SARS-CoV-2 virus), influenza, and respiratory syncytial virus (RSV). The dashboard will be updated every Friday at 1pm.

Weekly Surveillance Key Points

- Overall acute respiratory illness activity level in Chicago remains Low. Flu activity remains Low,
 COVID-19 remains Minimal, and RSV has decreased from Low to Minimal.
- ED visits, hospitalizations, and ICU admissions for flu, RSV, and COVID-19 remain low across all age groups.
- The percentage of specimens that tested positive for flu continues to decline as we approach the
 end of the influenza surveillance season. Of all influenza positive specimens last week, nearly all
 (90%) were associated with flu B. Test positivity for COVID-19 and RSV remain low.
- Compared to baseline levels, overall wastewater concentrations for RSV and COVID-19 remain
 low. Flu A has increased from low to moderate and flu B remains moderate. Even as clinical
 surveillance indicators (e.g., ED visits and hospitalizations) decrease, wastewater data may
 fluctuate for longer and decrease over an extended period of time because of the persistence of
 viral shedding from infected persons and/or in wastewater systems.
- The CDC has issued interim estimates of <u>2024-25 Seasonal Influenza Vaccine Effectiveness</u> (VE) and <u>2024-25 COVID-19 VE</u>, which suggest that this season's influenza and COVID-19 vaccines were effective in preventing medically-attended illness, including hospitalization, in the US.
- Flu and COVID-19 vaccine coverage across the city remain low. See CDPH's Mid-Season Seasonal Respiratory Vaccine Coverage Report for more details.
- For information on the national bird flu outbreak and local guidance see: the <u>CDPH H5N1 Bird Flu</u> webpage and <u>CDPH HAN: Flu Page</u> for clinicians.

Influenza-Associated ICU Hospitalizations - In Illinois, influenza-associated ICU hospitalizations are reportable as soon as possible but within three days. The graph below shows the weekly number of reported ICU hospitalizations for Chicago residents for previous two seasons. The table summarizes selected characteristics of reported cases for the current week and cumulative for the season.



	Week Ending		Since	
	May 3,	2025	September	29, 2024
Group	#	%	#	%
Citywide	1	100	531	100
Age				
0-4	0	0.0	38	7.2
5-17	0	0.0	37	7.0
18-24	0	0.0	15	2.8
25-49	0	0.0	94	17.7
50-64	0	0.0	108	20.3
65+	1	100	239	45.0
Gender				
Male	1	100	256	48.2
Female	0	0.0	275	51.8
Race-Ethnicity				
White Non-Latinx	1	100	125	23.5
Black Non-Latinx	0	0.0	244	46.0
Latinx	0	0.0	122	23.0
Asian Non-Latinx	0	0.0	23	4.3
Other Non-Latinx	0	0.0	11	2.1
Unknown-Race	0	0.0	6	1.1


Cinco

Respiratory Virus Laboratory Surveillance - Current Week and Cumulative The table below includes respiratory viral PCR tests performed by several hospital laboratories in Chicago as well as two commercial laboratories serving Chicago facilities. Reporting facilities represent nearly half of all acute care hospitals in the city. Data reported include Chicago and non-Chicago residents.

	Week Ending May 3, 2025		Since September 29, 2024	
Respiratory Pathogen	# Tested	% Positive	# Tested	% Positive
Influenza*	2,023	2.5	114,511	10.9
RSV*	1,227	0.7	92,581	5.5
SARS-CoV-2*	892	1.6	93,975	4.0
Parainfluenza	1,496	1.8	62,894	1.5
Rhinovirus/Enterovirus	602	17.9	33,709	12.9
Adenovirus	759	1.4	33,838	1.8
Human Metapneumovirus	767	6.3	34,285	1.6
Seasonal Coronaviruses [†]	1,488	0.8	53,129	2.7


^{*}Represents both dualplex and multiplex PCR data. All other data represents only multiplex panels that include the specified pathogens;† Four seasonal coronavirus strains include 229E, NL63, OC43, and HKU1.

Weekly number of specimens testing positive for influenza by type and subtype (graph) and the number of positive specimens by type and subtype for the current week and cumulative for the season (table).




	Week Ending		Since	
	May 3, 2025		September 29, 2024	
Type / Subtype	# Positive	%	# Positive	%
Influenza A	5	10.0	10,538	84.2
(H1N1)pdm09	1	20.0	1,175	11.2
H3N2	1	20.0	731	6.9
Subtyping not performed	3	60.0	8,632	81.9
Influenza B	45	90.0	1,975	15.8
Total Positive	50	100	12,513	100


Respiratory Virus Laboratory Surveillance - Seasonal Trends These graphs show seasonal trends of selected respiratory virus testing data presented in the previous table. Typical seasonal periods when activity tends to increase for influenza and RSV are indicated by shaded areas. Elevated test positivity outside of typical seasonal periods suggests atypical activity, and increased clinician awareness and testing may be warranted. Yearly data can also be used to compare the timing and intensity of viral activity, although changes in testing patterns also influence yearly trends, and data should be interpreted in the context of other surveillance indicators.




Emergency Department Illness Surveillance In Illinois, all 185 acute-care hospitals report emergency department visit data in near-real time to the Illinois Department of Public Health (IDPH). By tracking symptoms (or chief complaints) of patients in emergency departments, public health can promptly detect unusual levels of illness to determine whether a response is warranted.


Percent of emergency department visits attributed to **influenza-like illness (ILI)** for residents of Chicago zip codes based on chief complaint data.







Percent of emergency department visits attributed to **COVID-19 diagnoses** for residents of Chicago zip codes based on chief complaint data.






Percent of emergency department visits attributed to **respiratory syncytial virus (RSV)** diagnoses for residents of Chicago zip codes based on chief complaint data. Seasonal trends are displayed for children younger than 5 years old who are most impacted by RSV.






Outpatient Visit Illness Surveillance* Several outpatient clinics throughout Chicago participate in CDC's Influenza-like Illness Surveillance Network (<u>ILINet</u>) by reporting on a weekly basis the total number of outpatient clinic visits, and of those visits, the number with influenza-like illness (ILI). This graph shows the percent of medically-attended outpatient visits attributed to ILI as reported by ILINet facilities in Chicago.



Weekly Pediatric Admissions Emergency department visit data includes information on whether the visit resulted in a hospital admission at any time during the course of the clinical encounter. The syndromes or disease associated with the hospitalization are based on chief complaint and discharge diagnosis codes and no not necessarily represent lab-confirmed cases. The chart below represents hospital admissions among children <18 years-old at Chicago hospitals due to acute respiratory illnesses.



National and State Respiratory Virus Surveillance

The Centers for Disease Control and Prevention's FluView report provides national updates and trends related to influenza activity across the United States, and the National Respiratory and Enteric Virus Surveillance System (NREVSS) is a voluntary laboratory-based system that monitors temporal and geographic circulation patterns of several respiratory viruses in the U.S. The Respiratory Syncytial Virus (RSV) Hospitalization Surveillance Network (RSV-NET) is a CDC population-based surveillance system that collects data on severe RSV hospitalizations, including those resulting in ICU admission or death, among children and adults. The Respiratory Virus Hospitalization Surveillance Network (RESP-NET) monitors laboratory-confirmed hospitalizations associated with influenza, COVID-19, and respiratory syncytial virus (RSV) among children and adults. The Illinois and Suburban Cook County influenza surveillance reports are also available online. Current and archived issues of the Chicago Influenza and Respiratory Virus Surveillance Report can be found on the CDPH website Historical and Seasonal Summary Reports.