


### Chicago: Monkeypox (MPV) Update

July 22, 2022

## 2022 MPV global outbreak: 15,848 Total Confirmed Cases identified in 72 countries






Source: CDC, Data as of 7/21/2022

### In the U.S., 2,593 Total Confirmed MPV Cases identified









## While the overall risk of MPV remains low, see a healthcare provider if you develop symptoms

- As of 7/21, a total of 197 Chicagoans have been diagnosed with monkeypox virus (MPV)
  - Most, but not all, have been in gay, bisexual, or other men who have sex with men (MSM)
  - Median age of 35 years (range: 22-66)
  - Spread occurs through tight-knit social networks

#### If you have a new or unexplained rash:

- Visit a healthcare provider or call 312-746-4835 to get connected to care
- Remind your provider that MPV is in the community
- Avoid sex or being intimate with anyone until you've been checked out

#### Visual Examples of Monkeypox Rash





Photo Credit: NHS England High Consequence Infectious Disease









- Testing capacity has ramped up and is now plentiful
  - Effective diagnostic testing capacity, up to 70,000-80,000 tests can be performed per week nationally
  - Now in public health and commercial labs
  - Swabs of lesions can be collected by any medical provider
  - Please get tested if you have symptoms!

#### Treatment is available

- FDA-approved antiviral medication (EU-IND tecovirimat -TPOXX)
- Higher-risk patients
- Prescribed through health care providers; still significant paperwork





### MPV Vaccine

- A vaccine exists in limited quantities in the Strategic National Stockpile and the U.S. is acquiring more:
  - FDA-approved effective vaccines against MPV (JYNNEOS) help protect individuals who have been exposed to someone with MPV
  - Chicago has received and already distributed about 5,000 vaccine doses to date
  - 15,000 additional doses arriving from the federal government
  - State has also redirected some of its federal vaccine supply to Chicago, given epicenter of outbreak is here


### Ring Vaccination Strategy and Goal: Vaccinate All Contacts of Cases




Vaccinated

Unvaccinated











## YOU ARE ELIGIBLE FOR A MPV VACCINE IF YOU:

Had close physical contact with someone or an intimate partner who was diagnosed with MPV.

OR

2. Are a gay, bisexual or other (cis or trans) man who has sex with men.



Has intimate or sexual contact with other men in a social or sexual venue.

Has given or received money or other goods or services in exchange for sex.

Has intimate or sexual contact with multiple or anonymous partners.







**ADDITIONAL** 







At this time, MPV vaccine is NOT recommended for the general public, including same gender loving men, without other risk factors. As vaccine supply increases, guidance may evolve.



# FOR MORE INFO, VISIT ... CHI.GOV/MONKEYPOX



# Chicago is prioritizing first doses of JYNNEOS™ vaccine at this time

- At present, the top goal of immunization is outbreak control
- The 15,000 additional doses expected in Chicago will be used primarily as first doses to offer protection to more at-risk individuals
- Second doses will still be offered to immunocompromised individuals and to known contacts of MPV cases
- A second dose will be offered to others when vaccine supply improves
  - Biggest increase in protection is after the first dose
  - Delays in receiving the second dose do not affect the second dose immune response; 2<sup>nd</sup> dose vaccine labeled for "at least" 28 days later
  - Same approach in other countries