

**FIGURE 3. Catch-up immunization schedule for persons aged 4 months through 18 years who start late or who are more than 1 month behind —United States • 2012**

The figure below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age. **Always use this table in conjunction with the accompanying childhood and adolescent immunization schedules (Figures 1 and 2) and their respective footnotes.**

Persons aged 4 months through 6 years					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to dose 2	Dose 2 to dose 3	Dose 3 to dose 4	Dose 4 to dose 5
Hepatitis B	Birth	4 weeks	8 weeks and at least 16 weeks after first dose; minimum age for the final dose is 24 weeks		
Rotavirus <sup>1</sup>	6 weeks	4 weeks	4 weeks <sup>1</sup>		
Diphtheria, tetanus, pertussis <sup>2</sup>	6 weeks	4 weeks	4 weeks	6 months	6 months <sup>2</sup>
<i>Haemophilus influenzae</i> type b <sup>3</sup>	6 weeks	4 weeks if first dose administered at younger than age 12 months 8 weeks (as final dose) if first dose administered at age 12–14 months No further doses needed if first dose administered at age 15 months or older	4 weeks <sup>3</sup> if current age is younger than 12 months 8 weeks (as final dose) <sup>3</sup> if current age is 12 months or older and first dose administered at younger than age 12 months and second dose administered at younger than 15 months No further doses needed if previous dose administered at age 15 months or older	8 weeks (as final dose) This dose only necessary for children aged 12 months through 59 months who received 3 doses before age 12 months	
Pneumococcal <sup>4</sup>	6 weeks	4 weeks if first dose administered at younger than age 12 months 8 weeks (as final dose for healthy children) if first dose administered at age 12 months or older or current age 24 through 59 months No further doses needed for healthy children if first dose administered at age 24 months or older	4 weeks if current age is younger than 12 months 8 weeks (as final dose for healthy children) if current age is 12 months or older No further doses needed for healthy children if previous dose administered at age 24 months or older	8 weeks (as final dose) This dose only necessary for children aged 12 months through 59 months who received 3 doses before age 12 months or for children at high risk who received 3 doses at any age	
Inactivated poliovirus <sup>5</sup>	6 weeks	4 weeks	4 weeks	6 months <sup>5</sup> minimum age 4 years for final dose	
Meningococcal <sup>6</sup>	9 months	8 weeks <sup>6</sup>			
Measles, mumps, rubella <sup>7</sup>	12 months	4 weeks			
Varicella <sup>8</sup>	12 months	3 months			
Hepatitis A	12 months	6 months			
Persons aged 7 through 18 years					
Tetanus, diphtheria/ tetanus, diphtheria, pertussis <sup>9</sup>	7 years <sup>9</sup>	4 weeks	4 weeks if first dose administered at younger than age 12 months 6 months if first dose administered at 12 months or older	6 months if first dose administered at younger than age 12 months	
Human papillomavirus <sup>10</sup>	9 years	Routine dosing intervals are recommended <sup>10</sup>			
Hepatitis A	12 months	6 months			
Hepatitis B	Birth	4 weeks	8 weeks (and at least 16 weeks after first dose)		
Inactivated poliovirus <sup>5</sup>	6 weeks	4 weeks	4 weeks <sup>5</sup>	6 months <sup>5</sup>	
Meningococcal <sup>6</sup>	9 months	8 weeks <sup>6</sup>			
Measles, mumps, rubella <sup>7</sup>	12 months	4 weeks			
Varicella <sup>8</sup>	12 months	3 months if person is younger than age 13 years 4 weeks if person is aged 13 years or older			

- Rotavirus (RV) vaccines (RV-1 [Rotarix] and RV-5 [Rota Teq]).**
  - The maximum age for the first dose in the series is 14 weeks, 6 days; and 8 months, 0 days for the final dose in the series. Vaccination should not be initiated for infants aged 15 weeks, 0 days or older.
  - If RV-1 was administered for the first and second doses, a third dose is not indicated.
- Diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine.**
  - The fifth dose is not necessary if the fourth dose was administered at age 4 years or older.
- Haemophilus influenzae type b (Hib) conjugate vaccine.**
  - Hib vaccine should be considered for unvaccinated persons aged 5 years or older who have sickle cell disease, leukemia, human immunodeficiency virus (HIV) infection, or anatomic/functional asplenia.
  - If the first 2 doses were PRP-OMP (PedvaxHIB or Comvax) and were administered at age 11 months or younger, the third (and final) dose should be administered at age 12 through 15 months and at least 8 weeks after the second dose.
  - If the first dose was administered at age 7 through 11 months, administer the second dose at least 4 weeks later and a final dose at age 12 through 15 months.
- Pneumococcal vaccines.** (Minimum age: 6 weeks for pneumococcal conjugate vaccine [PCV]; 2 years for pneumococcal polysaccharide vaccine [PPSV])
  - For children aged 24 through 71 months with underlying medical conditions, administer 1 dose of PCV if 3 doses of PCV were received previously, or administer 2 doses of PCV at least 8 weeks apart if fewer than 3 doses of PCV were received previously.
  - A single dose of PCV may be administered to certain children aged 6 through 18 years with underlying medical conditions. See age-specific schedules for details.
  - Administer PPSV to children aged 2 years or older with certain underlying medical conditions. See *MMWR* 2010;59(No. RR-11), available at <http://www.cdc.gov/mmwr/pdf/rrr5911.pdf>.
- Inactivated poliovirus vaccine (IPV).**
  - A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose.
  - In the first 6 months of life, minimum age and minimum intervals are only recommended if the person is at risk for imminent exposure to circulating poliovirus (i.e., travel to a polio-endemic region or during an outbreak).
  - IPV is not routinely recommended for U.S. residents aged 18 years or older.
- Meningococcal conjugate vaccines, quadrivalent (MCV4).** (Minimum age: 9 months for Menactra [MCV4-D]; 2 years for Menveo [MCV4-CRM])
  - See Figure 1 ("Recommended immunization schedule for persons aged 0 through 6 years") and Figure 2 ("Recommended immunization schedule for persons aged 7 through 18 years") for further guidance.
- Measles, mumps, and rubella (MMR) vaccine.**
  - Administer the second dose routinely at age 4 through 6 years.
- Varicella (VAR) vaccine.**
  - Administer the second dose routinely at age 4 through 6 years. If the second dose was administered at least 4 weeks after the first dose, it can be accepted as valid.
- Tetanus and diphtheria toxoids (Td) and tetanus and diphtheria toxoids and acellular pertussis (Tdap) vaccines.**
  - For children aged 7 through 10 years who are not fully immunized with the childhood DTaP vaccine series, Tdap vaccine should be substituted for a single dose of Td vaccine in the catch-up series; if additional doses are needed, use Td vaccine. For these children, an adolescent Tdap vaccine dose should not be given.
  - An inadvertent dose of DTaP vaccine administered to children aged 7 through 10 years can count as part of the catch-up series. This dose can count as the adolescent Tdap dose, or the child can later receive a Tdap booster dose at age 11–12 years.
- Human papillomavirus (HPV) vaccines (HPV4 [Gardasil] and HPV2 [Cervarix]).**
  - Administer the vaccine series to females (either HPV2 or HPV4) and males (HPV4) at age 13 through 18 years if patient is not previously vaccinated.
  - Use recommended routine dosing intervals for vaccine series catch-up; see Figure 2 ("Recommended immunization schedule for persons aged 7 through 18 years").

Clinically significant adverse events that follow vaccination should be reported to the Vaccine Adverse Event Reporting System (VAERS) online (<http://www.vaers.hhs.gov>) or by telephone (800-822-7967). Suspected cases of vaccine-preventable diseases should be reported to the state or local health department. Additional information, including precautions and contraindications for vaccination, is available from CDC online (<http://www.cdc.gov/vaccines>) or by telephone (800-CDC-INFO [800-232-4636]).